



## Illinois Wesleyan University Digital Commons @ IWU

---

John Wesley Powell Student Research  
Conference

2003, 14th Annual JWP Conference

---

Apr 12th, 9:00 AM - 10:00 AM

# Synthesis of Hexamolybdate-Polyether Diamine Complexes

David Aggen

*Illinois Wesleyan University*

Rebecca Roesner, Faculty Advisor

*Illinois Wesleyan University*

Follow this and additional works at: <http://digitalcommons.iwu.edu/jwprc>

---

David Aggen and Rebecca Roesner, Faculty Advisor, "Synthesis of Hexamolybdate-Polyether Diamine Complexes" (April 12, 2003). *John Wesley Powell Student Research Conference*. Paper 1.  
<http://digitalcommons.iwu.edu/jwprc/2003/posters/1>

This Event is brought to you for free and open access by The Ames Library, the Andrew W. Mellon Center for Curricular and Faculty Development, the Office of the Provost and the Office of the President. It has been accepted for inclusion in Digital Commons @ IWU by the faculty at Illinois Wesleyan University. For more information, please contact [digitalcommons@iwu.edu](mailto:digitalcommons@iwu.edu).

©Copyright is owned by the author of this document.


## Poster Presentation P1

## SYNTHESIS OF HEXAMOLYBDATE-POLYETHER DIAMINE COMPLEXES

David Aggen and Rebecca Roesner\*

Department of Chemistry, Illinois Wesleyan University

Polyoxometalates are a class of highly symmetric anion clusters that have applications in supramolecular chemistry. These polyoxometalate clusters may be able to act as “ends” which trap macrocyclic rings on linear carbon chains. Polyoxometalates have already been incorporated into polymers<sup>1</sup> and dendrimers.<sup>2</sup> Our research involves the synthesis of a dumbbell shaped molecule through reaction of two equivalents of tetrabutylammonium hexamolybdate ( $[\text{Bu}_4\text{N}]_4[\text{Mo}_6\text{O}_{19}]$ ) and 1 equivalent of a polyether diamine strap. The diamine was prepared from 4-acetamidophenol and 1,2 bis(2-chloroethoxy)ethane. Further reaction of the prepared diamine with the hexamolybdate ion  $[\text{Mo}_6\text{O}_{19}]^{2-}$  was facilitated using dicyclohexylcarbodiimide ( $\text{C}_{13}\text{H}_{22}\text{N}_2$ ) which acts as a catalyst and a dehydrating agent. Dicyclohexylurea ( $\text{C}_{13}\text{H}_{24}\text{N}_2\text{O}$ ) is formed as a byproduct.<sup>3</sup> Formation of the diamine-hexamolybdate adduct was verified by  $^1\text{H}$  NMR and FT IR. We hope to refine our synthetic procedure to achieve higher yields.



<sup>1</sup> Moore, Aaron R.; Kwen, Haidoo; Beatty, Alicia M.; Maata, Eric A. *Chem. Comm.* **2000**, 1793-1794.

<sup>2</sup> Zeng, Huadong; Newkome, George R.; Hill, Craig L. *Angew. Chem. Int. Ed.* **2000**, 39, No. 10, 1772-1774.